

FORM PTO-1390 (REV. 9-2001)		U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE		ATTORNEY'S DOCKET NUMBER	
<b>TRANSMITTAL LETTER TO THE UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US) CONCERNING A FILING UNDER 35 U.S.C. 371</b>				AL01056K	
				U.S. APPLICATION NO. (if known, see 37 CFR 1.5)	
				<b>10/088708</b>	
INTERNATIONAL APPLICATION NO. PCT/US00/25595		INTERNATIONAL FILING DATE 19 September 2000 (19.09.00)		PRIORITY DATE CLAIMED 21 September 1999 (21.09.99)	
TITLE OF INVENTION USE OF DESLORATADINE FOR TREATING ALLERGIC AND INFLAMMATORY CONDITIONS					
APPLICANT(S) FOR DO/EO/US Melton B. Affrime, Christopher R. Banfield, Samir K. Gupta, Desmond Padhi					
Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information					
1. <input checked="" type="checkbox"/> This is a <b>FIRST</b> submission of items concerning a filing under 35 U.S.C. 371. 2. <input type="checkbox"/> This is a <b>SECOND</b> or <b>SUBSEQUENT</b> submission of items concerning a filing under 35 U.S.C. 371. 3. <input type="checkbox"/> This is an express request to begin national examination procedures (35 U.S.C. 371(f)). The submission must include items (5), (6), (9) and (21) indicated below. 4. <input type="checkbox"/> The US has been elected by the expiration of 19 months from the priority date (Article 31). 5. <input checked="" type="checkbox"/> A copy of the International Application as filed (35 U.S.C. 371(c)(2)) a. <input type="checkbox"/> is attached hereto (required only if not communicated by the International Bureau). b. <input type="checkbox"/> has been communicated by the International Bureau. c. <input checked="" type="checkbox"/> is not required, as the application was filed in the United States Receiving Office (RO/US). 6. <input type="checkbox"/> An English language translation of the International Application as filed (35 U.S.C. 371(c)(2)). a. <input type="checkbox"/> is attached hereto. b. <input type="checkbox"/> has been previously submitted under 35 U.S.C. 154(d)(4). 7. <input checked="" type="checkbox"/> Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3)) a. <input type="checkbox"/> are attached hereto (required only if not communicated by the International Bureau). b. <input type="checkbox"/> have been communicated by the International Bureau. c. <input type="checkbox"/> have not been made; however, the time limit for making such amendments has NOT expired. d. <input checked="" type="checkbox"/> have not been made and will not be made. 8. <input type="checkbox"/> An English language translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)). 9. <input checked="" type="checkbox"/> An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)). Copy of Int'l Appln as filed is attached. 10. <input type="checkbox"/> An English language translation of the annexes of the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)). 					
Items 11 to 20 below concern document(s) or information included:					
11. <input type="checkbox"/> An Information Disclosure Statement under 37 CFR 1.97 and 1.98. 12. <input type="checkbox"/> An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included. 13. <input checked="" type="checkbox"/> A <b>FIRST</b> preliminary amendment. 14. <input type="checkbox"/> A <b>SECOND</b> or <b>SUBSEQUENT</b> preliminary amendment. 15. <input type="checkbox"/> A substitute specification. 16. <input type="checkbox"/> A change of power of attorney and/or address letter. 17. <input type="checkbox"/> A computer-readable form of the sequence listing in accordance with PCT Rule 13ter.2 and 35 U.S.C. 1821 - 1.825 18. <input type="checkbox"/> A second copy of the published international application under 35 U.S.C. 154(d)(4). 19. <input type="checkbox"/> A second copy of the English language translation of the international application under 35 U.S.C. 154(d)(4) 20. <input checked="" type="checkbox"/> Other items or information: Preliminary Examination Report					



Patent Case No. AL01056K

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re application of: Melton Affrime, et al. :  
 For: TREATING ALLERGIC AND :  
 INFLAMMATORY CONDITIONS : Examiner: To Be Assigned

Serial No.: To Be Assigned : Group Art Unit: To Be Assigned

Filed: March 21, 2002 :

---

ASSISTANT COMMISSIONER FOR PATENTS  
 Washington, D.C. 20231

Schering-Plough Corp  
 Kenilworth, New Jersey 07033-0530

ATTN: Box Patent Application

SIR:

PRELIMINARY AMENDMENT

This application is the U.S. National Phase application of International Application No. PCT/US00/25595, filed 19 September 2000 which claims priority to U.S. Serial No. 09/400,147, filed September 21, 1999 and now abandoned.

The above-listed Title of the Invention is the title of the International application as filed and is the title on the executed Declaration and Power of Attorney form. The International Searching Authority in the International Search Report (dated 10 September 2001) changed the title to read as follows: "USE OF DESLORATADINE FOR TREATING ALLERGIC AND INFLAMMATORY CONDITIONS". Applicants authorize amendment of International title to the above-listed title if this is deemed necessary.

Please amend the specification as follows:

On Page 1, on line below the Title, add

"CROSS-REFERENCE TO RELATED APPLICATION: This application is the U.S. National Phase Application of International Application No. PCT/US00/25595, filed 19 September 2000, and claims priority to U.S. Serial No. 09/400,147, filed 09/21/1999, and now abandoned."

**IN THE CLAIMS:**

Please cancel claims 1-7, and add the following claims:

- (8) A method of treating or preventing allergic and inflammatory conditions of the skin or airway passages in a human in need of such treating or preventing while avoiding a food effect associated with other non-sedating antihistamines which comprises orally administering to said human under feed or fasted conditions an amount of desloratadine effective for such treating or preventing while avoiding a food effect on the bioavailability of desloratadine.
- (9) The method of claim 8 wherein the amount of desloratadine is about 2.5 mg/day to about 45 mg/day.
- (10) The method of claim 8 wherein the amount of desloratadine is about 2.5 mg/day.
- (11) The method of claim 8 wherein the amount of desloratadine is about 5 mg/day to about 10 mg/day.
- (12) The method of claim 8 wherein the amount of desloratadine is about 5 mg/day.
- (13) The method of claim 8 wherein the desloratadine is administered in a tablet formulation.
- (14) The method of claim 8 wherein the desloratadine is administered in a syrup formulation.
- (15) The method of claim 8 wherein the allergic reaction is season allergic rhinitis, perennial allergic rhinitis, atopic dermatitis, urticaria or allergic asthma.

- (16) A method of treating or preventing allergic and inflammatory conditions of the skin or airway passages in a human in need of such treating or preventing which comprises orally administering to said human an amount of desloratadine effective for such treating or preventing, while obtaining substantially the same bioavailability of desloratadine under feed or fasted conditions.
- (17) The method of claim 16 wherein the amount of desloratadine is about 2.5 mg/day to about 45 mg/day.
- (18) The method of claim 16 wherein the amount of desloratadine is about 2.5 mg/day.
- (19) The method of claim 16 wherein the amount of desloratadine is about 5 mg/day to about 10 mg/day.
- (20) The method of claim 16 wherein the amount of desloratadine is about 5 mg/day.
- (21) The method of claim 16 wherein the desloratadine is administered in a tablet formulation.
- (22) The method of claim 16 wherein the desloratadine is administered in a syrup formulation.
- (23) The method of claim 16 wherein the allergic reaction is season allergic rhinitis, pemnnial allergic rhinitis, atopic dermatitis, urticaria or allergic asthma.
- (24) A method of treating or preventing seasonal or perennial allergic rhinitis in a human in need of such treating or preventing while avoiding a food effect associated with non-sedating antihistamines which comprises orally administering to said human under feed or fasted conditions an amount of desloratadine effective for such treating or preventing while avoiding a food effect on the bioavailability of desloratadine.

- (25) The method of claim 24 wherein the amount of desloratadine is in the range of about 2.5 mg/day to about 45 mg/day.
- (26) The method of claim 24 wherein the amount of desloratadine is about 5 mg/day to about 15 mg/day.
- (27) The method of claim 24 wherein the amount of desloratadine is about 2.5 mg/day.
- (28) The method of claim 24 wherein the amount of desloratadine is about 5 mg/day.
- (29) The method of claim 24 wherein the patient is suffering from seasonal allergic rhinitis.
- (30) The method of claim 24 wherein the patient is suffering from perennial allergic rhinitis.
- (31) The method of claim 24 wherein the desloratadine is administered in a tablet formulation.
- (32) The method of claim 24 wherein the desloratadine is administered in a syrup formulation.
- (33) A method of treating or preventing seasonal or perennial allergic rhinitis in a human in need of such treating or preventing a which comprises orally administering to said human an amount of desloratadine effective for such treating or preventing, while obtaining substantially the same bioavailability of desloratadine under feed or fasted conditions.
- (34) The method of claim 33 wherein the amount of desloratadine is in the range of about 2.5 mg/day to about 45 mg/day.

- (35) The method of claim 33 wherein the amount of desloratadine is about 5 mg/day to about 15 mg/day.
- (36) The method of claim 33 wherein the amount of desloratadine is about 2.5 mg/day.
- (37) The method of claim 33 wherein the amount of desloratadine is about 5 mg/day.
- (38) The method of claim 33 wherein the patient is suffering from seasonal allergic rhinitis.
- (39) The method of claim 33 wherein the patient is suffering from perennial allergic rhinitis.
- (40) The method of claim 33 wherein the desloratadine is administered in a tablet formulation.
- (41) The method of claim 33 wherein the desloratadine is administered in a syrup formulation.
- (42) A method of treating or preventing atopic dermatitis or urticaria in a human in need of such treating or preventing while avoiding a food effect associated with non-sedating antihistamines which comprises orally administering to said human under feed or fasted conditions an amount of desloratadine effective for such treating or preventing while avoiding a food effect on the bioavailability of desloratadine.
- (43) The method of claim 42 wherein the amount of desloratadine is about 2.5 mg/day.
- (44) The method of claim 42 wherein the amount of desloratadine is about 5 mg/day to about 15 mg/day.
- (45) The method of claim 42 wherein the amount of desloratadine is about 5 mg/day to about 10 mg/day.
- (46) The method of claim 42 wherein the amount of desloratadine is about 5 mg/day.
- (47) The method of claim 42 wherein the patient is suffering from atopic dermatitis.
- (48) The method of claim 42 wherein the patient is suffering from urticaria.

(49) A method of treating or preventing atopic dermatitis or urticaria in a human in need of such treating or preventing a which comprises orally administering to said human an amount of desloratadine effective for such treating or preventing, while obtaining substantially the same bioavailability of desloratadine under feed or fasted conditions.

(50) The method of claim 49 wherein the amount of desloratadine is in the range of about 2.5 mg/day to about 45 mg/day.

(51) The method of claim 49 wherein the amount of desloratadine is about 2.5 mg/day to about 45 mg/day.

(52) The method of claim 49 wherein the amount of desloratadine is about 2.5 mg/day.

(53) The method of claim 49 wherein the amount of desloratadine is about 5 mg/day to about 10 mg/day.

(54) The method of claim 49 wherein the amount of desloratadine is about 5 mg/day.

(55) The method of claim 49 wherein the desloratadine is administered in a tablet formulation.

(56) The method of claim 49 wherein the desloratadine is administered in a syrup formulation.

(57) The method of claim 49 wherein the patient is suffering from atopic dermatitis.

(58) The method of claim 49 wherein the patient is suffering from urticaria.



**REMARKS**


Basis for the newly added claims 8-58 is found in claims 1-7 in the International Application and claims 1-13, as originally filed in the U.S. Serial No. 09/400,147, filed 09/22/1999, and in the specification, for example, in the Summary of the Invention, page 2, line 16-28, on page 2, line 31 to page 3, line 17, page 4, lines 10-29, page 12, line 1 to page 13, line 17, page 17, line 25 to page 18, line 12, and page 21, line 14 to page 22, line 5.

Newly added claims 8-58 are listed on Appendix I

No new matter will be added by entry of these claims.

Respectfully submitted

SCHERING-PLOUGH CORPORATION

A handwritten signature in black ink, appearing to read "Thomas D. Hoffman", with a stylized circular flourish at the end.

Thomas D. Hoffman

**APPENDIX I**

Please cancel claims 1-7, and add the following claims:

- (8) A method of treating or preventing allergic and inflammatory conditions of the skin or airway passages in a human in need of such treating or preventing while avoiding a food effect associated with other non-sedating antihistamines which comprises orally administering to said human under feed or fasted conditions an amount of desloratadine effective for such treating or preventing while avoiding a food effect on the bioavailability of desloratadine.
- (9) The method of claim 8 wherein the amount of desloratadine is about 2.5 mg/day to about 45 mg/day.
- (10) The method of claim 8 wherein the amount of desloratadine is about 2.5 mg/day.
- (11) The method of claim 8 wherein the amount of desloratadine is about 5 mg/day to about 10 mg/day.
- (12) The method of claim 8 wherein the amount of desloratadine is about 5 mg/day.
- (13) The method of claim 8 wherein the desloratadine is administered in a tablet formulation.
- (14) The method of claim 8 wherein the desloratadine is administered in a syrup formulation.
- (15) The method of claim 8 wherein the allergic reaction is season allergic rhinitis, pernninal allergic rhinitis, atopic dermatitis, urticaria or allergic asthma.
- (16) A method of treating or preventing allergic and inflammatory conditions of the skin or airway passages in a human in need of such treating or preventing a which comprises orally administering to said human an amount of desloratadine effective for such treating

or preventing, while obtaining substantially the same bioavailability of desloratadine under feed or fasted conditions.

(17) The method of claim 16 wherein the amount of desloratadine is about 2.5 mg/day to about 45 mg/day.

(18) The method of claim 16 wherein the amount of desloratadine is about 2.5 mg/day.

(19) The method of claim 16 wherein the amount of desloratadine is about 5 mg/day to about 10 mg/day.

(20) The method of claim 16 wherein the amount of desloratadine is about 5 mg/day.

(21) The method of claim 16 wherein the desloratadine is administered in a tablet formulation.

(22) The method of claim 16 wherein the desloratadine is administered in a syrup formulation.

(23) The method of claim 16 wherein the allergic reaction is season allergic rhinitis, perennial allergic rhinitis, atopic dermatitis, urticaria or allergic asthma.

(24) A method of treating or preventing seasonal or perennial allergic rhinitis in a human in need of such treating or preventing while avoiding a food effect associated with non-sedating antihistamines which comprises orally administering to said human under feed or fasted conditions an amount of desloratadine effective for such treating or preventing while avoiding a food effect on the bioavailability of desloratadine.

(25) The method of claim 24 wherein the amount of desloratadine is in the range of about 2.5 mg/day to about 45 mg/day.

(26) The method of claim 24 wherein the amount of desloratadine is about 5 mg/day to about 15 mg/day.

(27) The method of claim 24 wherein the amount of desloratadine is about 2.5 mg/day.

- (28) The method of claim 24 wherein the amount of desloratadine is about 5 mg/day.
- (29) The method of claim 24 wherein the patient is suffering from seasonal allergic rhinitis.
- (30) The method of claim 24 wherein the patient is suffering from perennial allergic rhinitis.
- (31) The method of claim 24 wherein the desloratadine is administered in a tablet formulation.
- (32) The method of claim 24 wherein the desloratadine is administered in a syrup formulation.
- (33) A method of treating or preventing seasonal or perennial allergic rhinitis in a human in need of such treating or preventing a which comprises orally administering to said human an amount of desloratadine effective for such treating or preventing, while obtaining substantially the same bioavailability of desloratadine under feed or fasted conditions.
- (34) The method of claim 33 wherein the amount of desloratadine is in the range of about 2.5 mg/day to about 45 mg/day.
- (35) The method of claim 33 wherein the amount of desloratadine is about 5 mg/day to about 15 mg/day.
- (36) The method of claim 33 wherein the amount of desloratadine is about 2.5 mg/day.
- (37) The method of claim 33 wherein the amount of desloratadine is about 5 mg/day.
- (38) The method of claim 33 wherein the patient is suffering from seasonal allergic rhinitis.
- (39) The method of claim 33 wherein the patient is suffering from perennial allergic rhinitis.

- (40) The method of claim 33 wherein the desloratadine is administered in a tablet formulation.
- (41) The method of claim 33 wherein the desloratadine is administered in a syrup formulation.
- (42) A method of treating or preventing atopic dermatitis or urticaria in a human in need of such treating or preventing while avoiding a food effect associated with non-sedating antihistamines which comprises orally administering to said human under feed or fasted conditions an amount of desloratadine effective for such treating or preventing while avoiding a food effect on the bioavailability of desloratadine.
- (43) The method of claim 42 wherein the amount of desloratadine is about 2.5 mg/day.
- (44) The method of claim 42 wherein the amount of desloratadine is about 5 mg/day to about 15 mg/day.
- (45) The method of claim 42 wherein the amount of desloratadine is about 5 mg/day to about 10 mg/day.
- (46) The method of claim 42 wherein the amount of desloratadine is about 5 mg/day.
- (47) The method of claim 42 wherein the patient is suffering from atopic dermatitis.
- (48) The method of claim 42 wherein the patient is suffering from urticaria.
- (49) A method of treating or preventing atopic dermatitis or urticaria in a human in need of such treating or preventing which comprises orally administering to said human an amount of desloratadine effective for such treating or preventing, while obtaining substantially the same bioavailability of desloratadine under feed or fasted conditions.
- (50) The method of claim 49 wherein the amount of desloratadine is in the range of about 2.5 mg/day to about 45 mg/day.
- (51) The method of claim 49 wherein the amount of desloratadine is about 2.5 mg/day to about 45 mg/day.
- (52) The method of claim 49 wherein the amount of desloratadine is about 2.5 mg/day.

- (53) The method of claim 49 wherein the amount of desloratadine is about 5 mg/day to about 10 mg/day.
- (54) The method of claim 49 wherein the amount of desloratadine is about 5 mg/day.
- (55) The method of claim 49 wherein the desloratadine is administered in a tablet formulation.
- (56) The method of claim 49 wherein the desloratadine is administered in a syrup formulation.
- (57) The method of claim 49 wherein the patient is suffering from atopic dermatitis.
- (58) The method of claim 49 wherein the patient is suffering from urticaria.

10/088708

TREATING ALLERGIC AND INFLAMMATORY CONDITIONSBACKGROUND OF THE INVENTION

This invention relates to treating and/or preventing allergic and inflammatory conditions in a human while avoiding a food effect associated with non-sedating antihistamines by administering an amount of desloratadine effective for such treating and/or preventing.

- 5 Loratadine is disclosed in U.S. Patent No. 4,282,233 as a non-sedating antihistamine useful for treating allergic reactions in animals including humans. Fexofenadine is disclosed in U.S. Patent Nos. 4,254,129 and 5,275,693 as a non-sedating antihistamine useful for treating allergic disorders.

- 10 Food intake induces changes in the physiology of the gastrointestinal tract that may influence drug absorption and/or drug clearance. Physiological changes induced by food intake can result in, *inter alia*, delayed gastric emptying, stimulation of bile flow, changes in pH, and increase in splanchnic blood flow. Food intake can also alter luminal metabolism and physically or chemically interact with a drug substance. The effects of co-administration of meals with
- 15 drugs is generally maximal when the drug product is administered immediately after completion of a meal. Meals that are high in calories, fat and density are likely to provide the greatest effects on bioavailability. While loratadine is safe and efficacious, there is a food effect associated with its administration. After co-administration of meals with loratadine, the effects of loratadine are higher than
- 20 when loratadine is administered to a patient under fasted condition [See Physician's Desk Reference (PDR), 54<sup>th</sup> Edition, 2000, Medical Economics Co., Montvale, N.J., at page 2782]. Fexofenadine has an opposite food effect; the effects of fexofenadine are greater when administered to a patient under fasted conditions (See CPS, 33<sup>rd</sup> Edition, 1998 Canadian Pharmacists Association,
- 25 Ottawa, Canada, at page 57).

In the case of another orally-active antihistamine, cetirizine hydrochloride, food had no effect on the cetirizine exposure (AUC), but the T<sub>max</sub> was delayed by

1.7 hours and the Cmax was decreased by 23% in the presence of food (See PDR, 54<sup>th</sup> Edition, 2000, at page 2404).

Thus, the complex effects of food, and the physicochemical properties and formulation of antihistamines often make the effect of food intake upon the bioavailability of antihistamines unpredictable.

Desloratadine is disclosed in U.S. Patent No. 4,659,716 as a non-sedating antihistamine useful for treating allergic reactions in animals including humans. We are aware of no publication regarding the association of food intake upon administration of desloratadine.

There is a need for a clinically effective therapy to treat or prevent such allergic and inflammatory conditions of the skin and airway passages in a human while avoiding the food effect associated with non-sedating antihistamines, such as loratadine and fexofenadine as well as other antihistamines such as cetirizine hydrochloride.

### **SUMMARY OF THE INVENTION**

The present invention provides a method of treating and/or preventing allergic and inflammatory conditions of the skin or airway passages in a human in need of such treating and /or preventing while avoiding a food effect associated with non-sedating antihistamines which comprises administering an amount of desloratadine effective for such treating and/or preventing.

The present invention also provides a method of treating and/or preventing seasonal or perennial allergic rhinitis in a human while avoiding a food effect associated with non-sedating antihistamines which comprises administering an amount of desloratadine effective for such treating and/or preventing.

The present invention provides a method of treating and/or preventing atopic dermatitis or urticaria in a human in need of such while avoiding a food effect associated with non-sedating antihistamines which comprises administering an amount of desloratadine effective for such treating and/or preventing.

### **DETAILED DESCRIPTION OF INVENTION**

The phrase "allergic and inflammatory conditions of the skin or airway passages" is meant those allergic and inflammatory conditions and symptoms



found on the skin and in the upper and lower airway passages from the nose to the lungs. Typical allergic and inflammatory conditions of the skin or upper and lower airway passages include seasonal and perennial allergic rhinitis, non-allergic rhinitis, asthma including allergic and non-allergic asthma, sinusitis, colds  
5 [in combination with a NSAID, e.g., aspirin or ibuprofen, or acetaminophen (APAP) and/or a decongestant, e.g., pseudoephedrine], dermatitis, especially allergic and atopic dermatitis, and urticaria and symptomatic dermographism as well as retinopathy, and small vessel diseases, associated with diabetes mellitus.

The amount of desloratadine effective for treating or preventing allergic and  
10 inflammatory conditions of the skin or airway passages will vary with the age, sex, body weight and severity of the allergic and inflammatory condition of the patient. Typically, the amount of desloratadine effective for treating or preventing such allergic and inflammatory conditions is in the range of about 2.5 mg/day to about 45 mg/day, preferably about 2.5 mg/day to about 20 mg/day, or about 5.0 mg/day  
15 to about 15 mg/day, or about 5.0 mg/day to about 10 mg/day, more preferably about 5.0 mg/day to about 7.5 mg/day, and most preferably about 5.0 mg/day in single or divided doses, or a single dose of 5.0 mg/day.

Desloratadine is a non-sedating long acting histamine antagonist with potent selective peripheral H1-receptor antagonist activity. Following oral  
20 administration, loratadine is rapidly metabolized to descarboethoxyloratadine or desloratadine, a pharmacologically active metabolite. *In vitro* and *in vivo* animal pharmacology studies have been conducted to assess various pharmacodynamic effects of desloratadine and loratadine. In assessing antihistamine activity in mice (comparison of ED<sub>50</sub> value), desloratadine was relatively free of producing  
25 alterations in behavior alterations in behavior, neurologic or autonomic function. The potential for desloratadine or loratadine to occupy brain H1-receptors was assessed in guinea pigs following i.p. administration and results suggest poor access to central histamine receptors for desloratadine or loratadine. *In vivo* studies also suggest that an inhibitory effect of desloratadine on allergic  
30 bronchospasm and cough can also be expected.

The clinical efficacy and safety of desloratadine has been documented in over 3,200 seasonal allergic rhinitis patients in 4 double-blinded, randomized clinical trials. The results of these clinical studies demonstrated the efficacy of

desloratadine in the treatment of adult and adolescent patients with seasonal rhinitis.

Efficacy endpoints in all the studies were Total Symptom Score, Total Nasal Symptom Score, Total Non-nasal Symptom Score, and Health Quality of

5 Life (HQOL) analysis in efficacy trials. Desloratadine (5 mg once daily) significantly reduced the total symptom scores (the sum of individual scores for rhinorrhea, sneezing, congestion/stuffiness, nasal itching, itchy/burning eyes, tearing, ocular redness, and itchy ears/palate). Desloratadine (5 mg) was significantly ( $p < 0.01$ ) more effective than placebo in reducing nasal symptoms.

10 An important efficacy endpoint analyzed in the desloratadine studies is the AM NOW total symptom score. This parameter measures the total symptom relief by the patient after 24 hours before taking the next day dose. Statistically significant ( $p < 0.05$ ) reductions were maintained for the full 24 hour dosing interval over the entire 5 mg to 20 mg dosage range

15 There were no significant differences in the effectiveness of desloratadine (over the entire 5 mg to 20 mg dosage range) across subgroups of patients defined by gender, age, or race. Desloratadine is particularly useful for the treatment and prevention of the nasal (stuffiness/congestion, rhinorrhea, nasal itching, sneezing) and non-nasal (itchy/burning eyes, tearing/watery eyes, redness  
20 of the eyes, itching of the ears/palate) symptoms of seasonal allergic rhinitis, including nasal congestion, in patients in need of such treating and/ or preventing.

Desloratadine is contraindicated in patients who are hypersensitive to this medication or to any of its ingredients.

#### **CLINICAL DESIGN FOR STUDY No. 1**

25 This open-label, two-way crossover study in 18 healthy subjects was designed to evaluate the effect of food on the bioavailability of Desloratadine in accordance with the FDA guidelines for evaluating the effect of food on the bioavailability of drugs. Subjects were randomized to receive single desloratadine 7.5 mg tablet under fasted conditions in one treatment period and immediately  
30 following a standardized high-fat, high-caloric breakfast meal in the other treatment period.



5 All subjects were confined to the study site until the 168-hour study related procedures were obtained. No strenuous physical activity was permitted, and the subjects were not allowed visitors while they were confined to the study site. A physician was present for all drug administrations and remained on the study site for at least four hours post-dose. A wash-out period of at least 7 days separated each period of the study.

10 Vital signs and ECGs were performed, and blood samples were collected at pre-specified times for safety and pharmacokinetic evaluations. Subjects were continually observed and questions throughout the study for possible occurrence of adverse events. Subjects were also instructed to report any unusual experiences or discomfort.

### **Overall Study Design: Discussion**

15 Since this study was conducted to determine the influence of food on the oral bioavailability (AUC and Cmax) of desloratadine, using the fasting state as reference, an open-label, randomized, two-way crossover design was used to meet the study objective.

### **Study Population/ Inclusion Criteria/ Exclusion Criteria**

20

#### **Inclusion Criteria:**

- Subjects were males or females between the ages of 18 and 45 years inclusive, and had a Body Mass Index (BMI) between 19-27.
  - Clinical laboratory tests (CBC, blood chemistries, urinalysis) were within normal limits or clinically acceptable to the Investigator/Sponsor .
  - Drug screen for drugs with a high potential for abuse were negative at screening and on admission to the study site.
  - Subjects were free of any clinically significant disease that required a physician's care and/or may have interfered with study evaluations, procedures or participation.
- 25
- 30

- Subject gave written informed consent (prior to any study-related procedures being performed) and were willing to adhere to restrictions and examination schedules.
- Subjects had a normal or clinically acceptable physical examination and ECG .

**Exclusion Criteria:**

- Subjects who had a history of any clinically significant local or systemic infectious disease within four weeks prior to initial treatment administration.
- Subjects who did not comply with the requirement that he or she should not have used any drugs (except acetaminophen) within 14 days prior to the study nor alcohol or xanthine-containing substances with 72 hours prior to study drug administration.
- Subjects who had participated in a clinical trial of any investigational drug within 30 days prior to the start of the study.
- Subjects who were, or were known to be former, narcotic addicts or alcoholics.
- Subjects who were positive for hepatitis B surface antigen or hepatitis C antibody.
- Subjects who were positive for HIV antibodies.
- Subjects who had a clinically significant history of food or drug allergy.
- Subjects who had a known allergy or intolerance to loratadine.
- Subjects who had donated blood within the preceding 30 days.
- Subjects who smoked, used tobacco products or used an adjunct to smoking cessation within the past 6 months (positive urine cotinine test).
- Females who were not surgically sterilized or were considering reversal of their surgical sterilization or were not at least 1 year post-menopausal.
- Females who had a positive urine pregnancy test at screening or on admission to the study site.
- Females who were lactating.

### Study Treatments

Subjects were confined to the study site at least 12 hours prior to each treatment administration. In the morning of Day 1 following a 10 hour overnight fast, each subject received one of the following treatments based on his/her subject number and the study period. The order of treatment administration was determined according to a computer-generated random code supplied to the Investigator by the Sponsor:

Treatment A: One desloratadine 7.5 mg tablet administered after a 10 hour fast.

Treatment B: One desloratadine 7.5 mg tablet administered immediately following a standardized high-fat, high-caloric breakfast.

Subjects randomized to receive the standardized high-fat, high-caloric breakfast (Treatment B) consumed the prescribed meal in a 20-minute period prior to drug administration and received desloratadine 7.5 mg tablet within 5 minutes after completing the breakfast.

Each dose was administered with 180 mL (6 fl oz) of non-carbonated room temperature water. The tablet was swallowed whole, not chewed or crushed. After dosing, the oral cavity was inspected to assure that the subject had swallowed the tablet. Subjects continued fasting (Treatment A) or did not eat again (Treatment B) until the 4- hour study procedures were completed, at which time lunch was served. Water was permitted throughout the fasting period except for 2 hours following treatment administration. The subjects remained awake and seated upright/ambulatory for 4 hours post-dose.. Subjects were under medical supervision throughout their confinement at the study site. Each treatment administration was separated by at least a 7 day washout period.

The desloratadine tablets were manufactured, packaged and supplied to the Investigator by Schering Corporation, Kenilworth, NJ, USA.

### Method of Treatment Assignment

This was a randomized, open-label, two-way crossover study. Upon confinement at the research center and after fulfilling all the study entry requirements, subjects were randomly assigned to received Treatment A (fasted condition) or Treatment B (fed condition) in one of the following two treatment-sequences according to a computer-generated random code supplied by Schering-Plough Research Institute.

AB

Or

BA

10

Subjects who withdrew or were removed from the study were to be replaced at the discretion of the Sponsor. Their replacement was to be numbered by using the original subject's assigned number plus an "R" (i.e., 1 replaced to 1R). The dosing regimen of the new subject was to be according to the original subject's dosing regimen.

15

### Selection of Doses & Selection and Timing of Dose for Each Subject

The desloratadine dose of 7.5 mg was selected because this was projected to be the clinical dose; similar response is expeted with other doses, e.g., 5 or 10 mg/day.

20

Each subject received a single 7.5 mg desloratadine tablet on two occasions.

### Prior and Concomitant Therapy

No other medications (investigational, prescription or OTC) except acetaminophen were taken by the subjects within 14 days of treatment initiation or during the course of the study without prior approval from the Principal Investigator or Sponsor unless it was a medical emergency. The use of any medications, including analgesics and over-the-counter medications that may have been used to treat adverse events, were to be recorded on the appropriate page of the case record form.

25

30

### Pharmacokinetics

Blood samples were collected for determination of the plasma pharmacokinetic profile of desloratadine. Fifteen milliliters (15mL) of blood were collected just prior to drug administration (0 hour) and at 0.5, 1, 1.5, 2, 3, 4, 6, 8, 12, 24, 36, 48, 72, 96, 120, 144 and 168 hours after dosing in both periods. All blood samples were collected into heparin-containing tubes at the specified times. The blood samples were centrifuged within 30 minutes after collection for 20 minutes at approximately 4°C and at approximately 3000 rpm. The plasma was separated and transferred into two separate appropriately labeled tubes, frozen to at least -20°C and maintained in the frozen state until assayed for desloratadine content.

The plasma concentration data for desloratadine (following administration of desloratadine 7.5 mg) was used to estimate the following pharmacokinetic parameters:

C<sub>max</sub> - maximum observed plasma concentration  
T<sub>max</sub> - time of observed maximum plasma concentration  
AUC(tf) - area under the plasma concentration vs time curve from time zero to the final measurable sampling time(tf)

AUC(l) - area under the plasma concentration vs time curve from time zero to infinity(l)

K - terminal phase rate constant

The major pharmacokinetic variables of interest were the plasma AUC(tf) and C<sub>max</sub>. All plasma samples were assayed for desloratadine concentration using a validated GLC/NPD method. The validation of the assay methods included documentation of its selectivity, limit of quantitation, linearity, precision and accuracy. The lower limit of quantitation (LOQ) of the assay was established at 0.1 ng/mL for desloratadine.



Plasma desloratadine concentration-time data were used to determine the pharmacokinetic parameters using model-independent methods. The maximum plasma concentration (C<sub>max</sub>) and time of maximum plasma concentration (T<sub>max</sub>) were the observed values. The terminal phase rate constant (K) was calculated as the negative of the slope of the log-linear terminal portion of the plasma concentration-time curve using linear regression. The terminal phase half-life (t<sub>1/2</sub>) was calculated as 0.693/K.

The area under the plasma concentration-time curve from time zero to the time of the final quantifiable sample [AUC(t<sub>f</sub>)] was calculated using the linear trapezoidal method and was extrapolated to infinity (I) according to the following equation:

$$AUC(I) = AUC(t_f) + C(t_f)K$$

where C(t<sub>f</sub>) is the estimated concentration at t<sub>f</sub>.

#### **Safety Measurements Assessed**

For safety evaluation, physical examinations, vital signs, electrocardiograms and clinical laboratory tests were conducted at screening and at the conclusion of the study (168 hours post-treatment). In addition, vital signs were monitored prior to treatment administration and daily during both treatment periods. Additional clinical laboratory tests and ECGs were obtained prior to dosing in each treatment period. The assessment, severity and relationship to treatment of adverse events were evaluated.

#### **SUMMARY CONCLUSIONS FOR Study No. 1:**

**RESULTS:** Study No. 1 was conducted as planned.

**Clinical Pharmacology:** The pharmacokinetics for the two-way cross-over study are presented in Tables I and II hereinbelow.

Table I

The Mean (%CV) pharmacokinetic parameters of desloratadine following oral, single-dose administration of 7.5 mg under fed and fasted conditions:			
	N	FED	FASTED
Parameter (Unit)		Mean (%CV) <sup>1</sup>	Mean (%CV)
Cmax(ng/mL)	18	3.53 (33)	3.30 (36)
AUC(tf)(ng.hr/mL)	18	73.8 (81)	77.5(92)
AUC(l)(ng.hr/mL)	17	62.5(40)	63.5(45)

1. %CV is percent coefficient of variation, which is a relative measure of variability.

See Steele and Torrie, "Principles and Procedures of Statistics", (1980) 2<sup>nd</sup>

5 Edition, McGraw-Hill, NY, at page 27.

TABLE II

The estimates of bioavailability (log-transformed) of desloratadine under fed condition relative to that after fasting		
Parameter	Point Estimate(%) <sup>c</sup>	90% Confidence Interval <sup>d</sup>
Cmax(ng/mL) <sup>a</sup>	108	99-118
AUC(tf)(ng.hr/mL) <sup>a</sup>	100	93-107
AUC(l)(ng.hr/mL) <sup>b</sup>	101	93-108

a: n=18

b: n=17

c: Ratio of the mean value for fed vs. fasted.

d:  $\alpha=0.05$ (two-tailed)

**Safety:** Blood pressure, pulse rate, respiratory rate, oral body temperature and electrocardiogram evaluations showed no consistent changes of clinical relevance and remained within the range observed for healthy subjects. Overall, 9 of 18 (50%) subjects reported treatment-emergent adverse events. The most frequently reported adverse event was headache. All reported adverse events were mild in severity except one such event that was reported as moderated. No subject discontinued participation in the study due to adverse events and no intervention was required to treat any adverse event.

#### **CONCLUSIONS FOR STUDY No. 1:**

- Single oral doses of desloratadine 7.5 mg administered under fed and fasted conditions were safe and well tolerated.
- Food intake had no effect on the oral bioavailability of desloratadine from the tablet formulation.

#### **CLINICAL DESIGN FOR STUDY No. 2**

This open-label, three-way crossover study in 30 healthy subjects was designed to evaluate the effect of food on the bioavailability of desloratadine syrup in accordance with the FDA guidelines for evaluating the effect of food on the bioavailability of drugs and to determine the bioequivalence of desloratadine between the tablet and syrup formulations. Subjects were randomized to receive one desloratadine 5.0 mg tablet under fasted conditions in a one treatment period or ten(10) mL of desloratadine syrup (0.5mg/mL) under fasted conditions in a second treatment period or ten(10) mL of desloratadine syrup(0.5mg/mL) immediately following a standardized high-fat, high-caloric breakfast meal in third treatment period.

#### **STUDY OBJECTIVE**

The objectives of Study No. 2 were to determine the bioequivalency of desloratadine between the tablet and syrup formulations and to evaluate the effect

of food on the bioavailability of desloratadine following administration of a syrup formulation.

# **OVERALL STUDY DESIGN AND PLAN:**

5 A total of 30 healthy subjects were enrolled and successfully completed this randomized, open-label, single-dose, three-way crossover study.

Subjects were screened within 3 weeks of dosing, and those who met the entry criteria were confined to the study center within 12 hours prior to each treatment (Day -1). Upon confinement, subjects had safety laboratory tests and  
10 electrocardiograms repeated. The following morning, after fasting for a minimum of 10 hours, subjects received one of the following treatments based on his/her subject number and the study period:

- |    |              |   |
|----|--------------|---|
| 15 | Treatment A: | One desloratadine 5 mg tablet administered after a 10-hour fast.  |
|    | Treatment B: | Ten (10) mL of desloratadine syrup (0.5 mg/mL) following a 10-hour fast.  |
| 20 | Treatment C: | Ten (10) mL of desloratadine syrup (0.5 mg/mL) administered immediately following a standardized high-fat and high-caloric breakfast. |

Subjects who were randomized to receive the standardized high-fat and high-caloric breakfast (Treatment C), consumed the prescribed meal in a  
25 20-minute period prior to drug administration and received the appropriate dose of desloratadine within 5 minutes after completing the breakfast.

The tablets were administered with 180 mL (6 fl oz) of noncarbonated room temperature water. The tablet was swallowed whole, not chewed or crushed. After dosing, the oral cavity was inspected to assure that the subject swallowed  
30 the tablet/syrup. For subjects randomized to Treatment B or Treatment C the

study medication was administered by having the volunteer drink the entire 10 mL of desloratadine syrup, followed by two 10 mL tap water rinses of the dose container (ie, oral syringe, etc.) to ensure complete dose intake. Subjects consumed the remaining 160 mL of water. Subjects continued fasting  
5 (Treatments A and B) or did not eat again (Treatment C) until the 4-hour study procedures were completed, at which time lunch was served. Water was permitted throughout the fasting period except for 2 hours following treatment. The subjects remained awake and seated upright/ambulatory for 4 hours post-dose.

10 All subjects were confined to the study site until the 120-hour study related procedures were obtained. No strenuous physical activity was permitted, and the subjects were not allowed visitors while they were confined to the study site. A physician was present for all drug administrations and remained on the study site for at least four hours post-dose. A washout period of at least 14 days separated  
15 each period of the study.

Vital signs and ECGs were performed and blood samples were collected at prespecified times for safety and pharmacokinetic evaluations. Subjects were continually observed and questioned throughout the study for possible occurrence of adverse events. Subjects were also instructed to report any unusual  
20 experiences or discomfort.

#### OVERALL STUDY DESIGN:

Since Study No. 2 was conducted to determine the bioequivalency of desloratadine between the tablet and syrup formulations and to evaluate the  
25 influence of food on the oral bioavailability (AUC and C<sub>max</sub>) of desloratadine of the syrup formulation an open-label, randomized, three-way crossover design was used to meet the study objective.

The Inclusion Criteria and Exclusion Criteria used for Study No. 2 were the same as those used for Study No. 1 except that:



Statistical comparisons of log-transformed Cmax and AUC(I) values for Treatment B versus Treatment A and Treatment C versus Treatment B were performed for both desloratadine and 3-OH desloratadine. Overall, the results suggested that there were no statistically significant differences in the bioavailabilities of desloratadine from the tablet and syrup formulations, and that

high-fat and high-caloric meals had no effect on the bioavailability of desloratadine from the syrup formulation.

The relative bioavailability and the 90% Confidence Intervals (CIs) for the log-transformed C<sub>max</sub> and AUC(I) for desloratadine and 3-OH desloratadine are presented in Table IV. The CIs of AUC(I) and C<sub>max</sub> of desloratadine and 3-OH desloratadine for Treatment B relative to Treatment A met the 80-125% bioequivalence guideline. This indicates that the tablet and syrup formulations were bioequivalent. The corresponding CIs for the syrup under fasted and fed conditions met the bioequivalency criteria for C<sub>max</sub> (70-143%) and AUC (80-125%). Thus, a high-fat and high-caloric meal had no effect on the bioavailability of desloratadine or 3-OH desloratadine from the syrup formulation. There was also no effect of food previously seen with the tablet formulation.



TABLE III

**Mean (%CV) Pharmacokinetic Parameters of Desloratadine and 3-OH Desloratadine in Healthy Adult Volunteers Following Single-Dose Oral Administration of Desloratadine Tablet and Syrup Formulations Under Either Fasted or Fed Condition (N=30)**

Pharmacokinetic Parameters			
Treatment	C <sub>max</sub> (ng/mL)	AUC(t <sub>f</sub> ) (ng·hr/mL)	AUC(l) (ng·hr/mL)
Desloratadine			
A	2.44 (41)	45.8 (44)	47.4 (45)
B	2.30 (51)	46.2 (71)	48.4 (74)
C	2.19 (62)	49.9 (90)	52.0 (93)
3-OH Desloratadine			
A	1.06 (34)	27.0 (25)	29.0 (24)
B	1.03 (38)	26.0 (28)	27.8 (28)
C	0.91 (38)	25.8 (31)	28.3 (31)

Where:

Treatment A=One 5 mg desloratadine tablet administered after a 10-hour fast.

Treatment B=Ten (10) mL desloratadine syrup (0.5 mg/mL) administered after a 10-hour fast.

Treatment C=Ten (10) mL desloratadine syrup (0.5 mg/mL) administered immediately following a standardized high-fat and high-caloric breakfast.

TABLE IV

**Estimates of Bioequivalence and the 90% Confidence Intervals for the Log-Transformed C<sub>max</sub> and AUC(I) for Desloratadine and 3-OH Desloratadine in Healthy Adult Volunteers Following Single-Dose Oral Administration of Desloratadine Tablet and Syrup Formulations Under Fasted or Fed Condition**

Formulations Compared		Relative Bioavailability (%)	Confidence Interval (%) <sup>a</sup>
<b>Desloratadine</b>			
Treatment B/Treatment A	AUC(I)	95.4	84-108
	C <sub>max</sub>	92.5	84-102
Treatment C/Treatment B	AUC(I)	104	92-118
	C <sub>max</sub>	94.1	85-104
<b>3-OH Desloratadine</b>			
Treatment B/Treatment A	AUC(I)	94.9	89-101
	C <sub>max</sub>	96.5	89-104
Treatment C/Treatment B	AUC(I)	101	95-108
	C <sub>max</sub>	87.2	81-94

a: Ninety percent confidence interval.

Where:

Treatment A=One 5 mg desloratadine tablet administered after a 10-hour fast.

Treatment B=Ten (10) mL desloratadine syrup (0.5 mg/mL) administered after a 10-hour fast.

Treatment C=Ten (10) mL desloratadine syrup (0.5 mg/mL) administered immediately following a standardized high-fat, high-caloric breakfast.

**SAFETY:** Overall, 14 of 30 subjects (47%) reported at least one adverse event ("AE") during the study. Eight of 30 (27%) subjects reported at least one AE during the fasted treatment period with the tablet, 4 of 30 (13%) subjects reported at least one AE during the fasted treatment period with the syrup and 7 of 30 (23%) subjects reported at least one AE during the fed treatment period with the syrup. The most common AE (regardless of association to treatment) was headache.

Six and 5 subjects reported headaches of mild and moderate severity, respectively. All headaches rated as moderate were treated with acetaminophen.

Ten of 30 subjects (33%) AEs were considered to be treatment-related treatment-emergent AEs. The most common treatment-related AEs were headache (10 of 30; 33%) and gastrointestinal disorders (2/30; 7%).

5 No serious nor unexpected AEs were reported. No subject discontinued participation in the study due to AEs.

There were no deaths or serious or significant adverse events. There were no clinically significant abnormal laboratory values. Blood pressure, pulse rate, electrocardiograms and oral body temperature evaluations showed no consistent changes of clinical relevance and remained within the range observed for healthy male and female subjects.

## OVERALL CONCLUSIONS FOR STUDY NO. 2:

15 The administration of desloratadine 5 mg, in either a tablet or syrup formulation, was found to be safe and well tolerated in this study. All adverse events were considered mild to moderate in severity and the incidence of adverse events was similar across treatments. No subject was discontinued from the study due to an adverse event.

20 In Study, the bioequivalence of the 5 mg desloratadine syrup formulation (0.5 mg/mL) was evaluated with the 5 mg desloratadine tablet formulation. Based on the 90% CIs for C<sub>max</sub> and AUC, the 2 formulations were found to be bioequivalent with respect to desloratadine as well as its major metabolite, 3-OH desloratadine. In addition, the relative bioavailability of the syrup under fed and fasted conditions was evaluated. No food effect was observed for the syrup as evidenced by the 90% CIs  
25 which also met the bioequivalence guidelines.

## CONCLUSIONS FOR STUDY NO. 2:

- The tablet and syrup desloratadine formulations were bioequivalent.
- High-fat and high-caloric meal had no effect on the bioavailability of  
30 desloratadine from the syrup formulation.

## CONCLUSIONS FOR STUDY NOs. 1 & 2:

Thus, food intake had no effect on the bioavailability of oral administrated desloratadine from the tablet or syrup formulations. Similar results are expected when the effect of food intake on the bioavailability of the rapidly-disintegrating desloratadine formulation is evaluated.

### GENERAL EXPERIMENTAL

U.S. Patent No. 4,659,716 discloses methods of making desloratadine, pharmaceutical compositions containing it and methods of using desloratadine and pharmaceutical compositions containing it to treat allergic reaction in mammals.

U.S. Patent No. 5,595,997 discloses pharmaceutical compositions containing desloratadine and methods of using desloratadine for treating and preventing various disease states, e.g., allergic rhinitis.

Desloratadine is available from Schering Corporation, Kenilworth, N.J.

The pharmaceutical compositions of desloratadine can be adapted for any mode of administration e.g., for oral, e.g. tablet, syrup or rapidly-disintegrating tablet, parenteral, e.g., subcutaneous ("SC"), intramuscular ("IM"), and intraperitoneal ("IP"), topical or vaginal administration or by inhalation (orally or intranasally). Preferably desloratadine is administered orally.

Such pharmaceutical compositions may be formulated by combining desloratadine or an equivalent amount of a pharmaceutically acceptable salt thereof with a suitable, inert, pharmaceutically acceptable carrier or diluent that may be either solid or liquid. Desloratadine may be converted into the pharmaceutically acceptable acid addition salts by admixing it with an equivalent amount of a pharmaceutically acceptable acid. Typically suitable pharmaceutically acceptable acids include the mineral acids, e.g., HNO<sub>3</sub>, H<sub>2</sub>SO<sub>4</sub>, H<sub>3</sub>PO<sub>4</sub>, HCl, HBr, organic acids, including, but not limited to, acetic, trifluoroacetic, propionic, lactic, maleic, succinic, tartaric, glucuronic and citric acids as well as alkyl or arylsulfonic acids, such as p-toluenesulfonic acid, 2-naphthalenesulfonic acid, or methanesulfonic acid. The preferred pharmaceutically acceptable salts are trifluoroacetate, tosylate, mesylate, and chloride. Desloratadine is more stable as

the free base than as an acid addition salt and the use of the desloratadine free base in pharmaceutical compositions of the present invention is more preferred.

Solid form preparations include powders, tablets, dispersible granules, capsules, cachets and suppositories. The powders and tablets may be comprised of from about 5 to about 95 percent active ingredient. Suitable solid carriers are known in the art, e.g. magnesium carbonate, magnesium stearate, talc, sugar or lactose. Tablets, powders, cachets and capsules can be used as solid dosage forms suitable for oral administration. Examples of pharmaceutically acceptable carriers and methods of manufacture for various compositions may be found in A. Gennaro (ed.), Remington's Pharmaceutical Sciences, 18th Edition, (1990), Mack Publishing Co., Easton, Pennsylvania.

Liquid form preparations include solutions, suspensions and emulsions. As an example may be mentioned water or water-propylene glycol solutions for parenteral injection. Solid form preparations may be converted into liquid preparations shortly before use for either oral or administration. Parenteral forms to be injected intravenously, intramuscularly or subcutaneously are usually in the form of sterile solutions and may contain tonicity agents (salts or glucose), and buffers. Opacifiers may be included in oral solutions, suspensions and emulsions. Liquid form preparations may also include solutions for intranasal administration.

Aerosol preparations suitable for inhalation may include solutions and solids in powder form, which may be in combination with a pharmaceutically acceptable carrier, such as an inert compressed gas, e.g., nitrogen.

Also included are solid form preparations which are intended to be converted, shortly before use, to liquid form preparations for either oral or parenteral administration. Such liquid forms include solutions, suspensions and emulsions.

The compounds of the invention may also be deliverable transdermally. The transdermal compositions can take the form of creams, lotions, aerosols and/or emulsions and can be included in a transdermal patch of the matrix or reservoir type as are conventional in the art for this purpose.

Preferably, the pharmaceutical preparation is in a unit dosage form. In such form, the preparation is subdivided into suitably sized unit doses containing

WO 01/21161

PCT/US00/25595

24

appropriate quantities of the active component, e.g., an effective amount to achieve the desired purpose.

What is claimed:

- (1) The use of desloratadine for the preparation of a medicament for treating and/or preventing allergic and inflammatory conditions of the skin or airway passages in a human in need of such treating and /or preventing while avoiding a food effect associated with non-sedating antihistamines wherein the medicament comprises an amount of desloratadine effective for such treating and/or preventing.
- (2) The use of desloratadine for the preparation of a medicament for treating and/or preventing seasonal or perennial allergic rhinitis in a human while avoiding a food effect associated with non-sedating antihistamines wherein the medicament comprises an amount of desloratadine effective for such treating and/or preventing.
- (3) The use of desloratadine for the preparation of a medicament for treating and/or preventing atopic dermatitis or urticaria in a human in need of such while avoiding a food effect associated with non-sedating antihistamines wherein the medicament comprises an amount of desloratadine effective for such treating and/or preventing.
- (4) The use of any preceding claim wherein the amount of desloratadine is about 2.5 mg/day to about 45 mg/day.
- (5) The use of any preceding claim wherein the amount of desloratadine is about 5 mg/day to about 15 mg/day.
- (6) The use of any preceding claim wherein the amount of desloratadine is about 5 mg/day to about 10 mg/day.
- (7) The use of any preceding claim wherein the amount of desloratadine is about 5 mg/day in single or divided doses.

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
29 March 2001 (29.03.2001)

PCT

(10) International Publication Number  
WO 01/21161 A2

(51) International Patent Classification: A61K 31/00

(74) Agent: HOFFMAN, Thomas, D.; Schering-Plough Corporation, Patent Department K-6-1-1990, Kenilworth, NJ 07033-0530 (US).

(21) International Application Number: PCT/US00/25595

(22) International Filing Date:  
19 September 2000 (19.09.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
09/400,147 21 September 1999 (21.09.1999) US

(71) Applicant (for all designated States except US): SCHERING CORPORATION [US/US]; 2000 Galloping Hill Road, Kenilworth, NJ 07033-0530 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, HR, HU, ID, IL, IN, IS, JP, KG, KR, KZ, LC, LK, LR, LT, LU, LV, MA, MD, MG, MK, MN, MX, MZ, NO, NZ, PL, PT, RO, RU, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, US, UZ, VN, YU, ZA.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (for US only): AFFRIME, Melton, B. [US/US]; 11 Whispering Way, Warren, NJ 07059 (US). BANFIELD, Christopher, R. [US/US]; 4 Robin Lane, High Bridge, NJ 08829 (US). GUPTA, Samir, K. [US/US]; 14 Dobson Road, East Brunswick, NJ 08816 (US). PADHI, Desmond [CA/US]; 2026 Warble Court, Thousand Oaks, CA 91320 (US).

Published:

— Without international search report and to be republished upon receipt of that report.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 01/21161 A2

(54) Title: TREATING ALLERGIC AND INFLAMMATORY CONDITIONS

(57) Abstract: The use of desloratadine for the preparation of a medicament for treating and/or preventing allergic and inflammatory conditions of the skin or upper and lower airway passages in a human while avoiding the food effect associated with non-sedating antihistamines, e.g., loratadine or fexofenadine is disclosed.



Please type e plus sign (+) inside this box → ☐

PTO/SB/01 (12-97)  
 Approved for use through 9/30/00. OMB 0651-0032  
 Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number.

<b>DECLARATION FOR UTILITY OR DESIGN PATENT APPLICATION (37 CFR 1.63)</b>  <input checked="" type="checkbox"/> Declaration Submitted with Initial Filing    OR <input type="checkbox"/> Declaration Submitted after Initial Filing (surcharge (37 CFR 1.16 (e)) required)	<b>Attorney Docket Number</b>	AL01056K
	<b>First Named Inventor</b>	Melton B. Affrime
	<b>COMPLETE IF KNOWN</b>	
	<b>Application Number</b>	/
	<b>Filing Date</b>	
	<b>Group Art Unit</b>	
	<b>Examiner Name</b>	

**As a below named inventor, I hereby declare that:**

My residence, post office address, and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

TREATING ALLERGIC AND INFLAMMATORY CONDITIONS

the specification of which (Title of the Invention)

☒ is attached hereto  
 OR  
☐ was filed on (MM/DD/YYYY) \_\_\_\_\_ as United States Application Number or PCT International Application Number \_\_\_\_\_ and was amended on (MM/DD/YYYY) \_\_\_\_\_ (if applicable).

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment specifically referred to above.

I acknowledge the duty to disclose information which is material to patentability as defined in 37 CFR 1.56.

I hereby claim foreign priority benefits under 35 U.S.C. 119(a)-(d) or 365(b) of any foreign application(s) for patent or inventor's certificate, or 365(e) of any PCT international application which designated at least one country other than the United States of America, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate, or of any PCT international application having a filing date before that of the application on which priority is claimed.

Prior Foreign Application Number(s)	Country	Foreign Filing Date (MM/DD/YYYY)	Priority Not Claimed	Certified Copy Attached?	
				YES	NO
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

☐ Additional foreign application numbers are listed on a supplemental priority data sheet PTO/SB/02B attached hereto:

I hereby claim the benefit under 35 U.S.C. 119(a) of any United States provisional application(s) listed below.

Application Number(s)	Filing Date (MM/DD/YYYY)

☐ Additional provisional application numbers are listed on a supplemental priority data sheet PTO/SB/02B attached hereto.

[Page 1 of 2]

**CERTIFICATE OF MAILING**

I hereby certify that this correspondence is being deposited with the United States Postal Service as first class mail in an envelope addressed to: Assistant Commissioner for Patents, Washington, D.C. 20231 on this date: \_\_\_\_\_

Typed or printed name: THOMAS D. HOFFMAN

Signature

Date

Express Mail Label No. \_\_\_\_\_

Date \_\_\_\_\_

Please type e plus sign (+) inside this box ☐

PTO/SB/01 (12-97)

Approved for use through 9/30/00. OMB 0651-0032

Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number.

**DECLARATION — Utility or Design Patent Application**

I hereby claim the benefit under 35 U.S.C. 120 of any United States application(s), or 365(c) of any PCT international application designating the United States of America, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT International application in the manner provided by the first paragraph of 35 U.S.C. 112, I acknowledge the duty to disclose information which is material to patentability as defined in 37 CFR 1.56 which became available between the filing date of the prior application and the national or PCT international filing date of this application.

U.S. Parent Application or PCT Parent Number	Parent Filing Date (MM/DD/YYYY)	Parent Patent Number (if applicable)
09/400,147	09/21/00	

☐ Additional U.S. or PCT international application numbers are listed on a supplemental priority date sheet PTO/SB/02B attached hereto.

As a named inventor, I hereby appoint the following registered practitioner(s) to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith: ☐ Customer Number ☐ OR ☒ Registered practitioner(s) name/registration number listed below

Name	Registration Number	Name	Registration Number
Thomas D. Hoffmann	28,221		

☒ Additional registered practitioner(s) named on supplemental Registered Practitioner Information sheet PTO/SB/02C attached hereto.

Direct all correspondence to: ☐ Customer Number or Bar Code Label ☐ 24265 OR ☒ Correspondence address below

Name	Thomas D. Hoffmann				
Address	Patent Dept. K-6-1, 1990				
Address	2000 Galloping Hill Road				
City	Kenilworth	State	NJ	ZIP	07033-0530
Country	USA	Telephone	(908) 298-5037	Fax	(908) 298-5388

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under 18 U.S.C. 1001 and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Name of Sole or First Inventor: ☐ A petition has been filed for this unsigned inventor

Given Name (first and middle (if any))	Family Name or Surname
Melton B.	Affrine

Inventor's Signature	Date
<i>Melton B. Affrine</i>	12/11/00

Residence: City	Warren	State	NJ	Country	U.S.A.	Citizenship	U.S.A.
-----------------	--------	-------	----	---------	--------	-------------	--------

Post Office Address	11 Whispering Way				
---------------------	-------------------	--	--	--	--

Post Office Address					
---------------------	--	--	--	--	--

City	Warren	State	NJ	ZIP	07059	Country	U.S.A.
------	--------	-------	----	-----	-------	---------	--------

☒ Additional inventors are being named on the 1 supplemental Additional Inventor(s) sheet(s) PTO/SB/02A attached hereto

Please type a plus sign (+) inside this box → ☐

PTO/SB/02A (3-97)

Approved for use through 9/30/98. OMB 0651-0032

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number.

Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE

**DECLARATION****ADDITIONAL INVENTOR(S)**

Supplemental Sheet

Page 1 of 1

**Name of Additional Joint Inventor, If any:**☐ A petition has been filed for this unsigned inventor

Given Name (first and middle (if any))

Family Name or Surname

Christopher R.

Banfield

Inventor's  
Signature*Christopher R Banfield*

Date

19 Oct, 2002

Residence: City

High Bridge

State

NJ

Country

U.S.A.

Citizenship

U.S.A. *USX*

Post Office Address

4 Robin Lane

Post Office Address

City

High Bridge

State

NJ

ZIP

08829

Country

U.S.A.

**Name of Additional Joint Inventor, if any:**☐ A petition has been filed for this unsigned inventor

Given Name (first and middle (if any))

Family Name or Surname

Samir K.

Gupta

Inventor's  
Signature*Samir K Gupta*

Date

19 Oct, 2002

Residence: City

East Brunswick

State

NJ

Country

U.S.A.

Citizenship

U.S.A. *USX*

Post Office Address

14 Dobson Road

Post Office Address

City

East Brunswick

State

NJ

ZIP

08816

Country

U.S.A.

**Name of Additional Joint Inventor, If any:**☐ A petition has been filed for this unsigned inventor

Given Name (first and middle (if any))

Family Name or Surname

Desmond

Padhi

Inventor's  
Signature*Desmond Padhi*

Date

16 Oct, 2002

Residence: City

Thousand Oaks

State

CA

Country

U.S.A.

Citizenship

Canada. *CAX*

Post Office Address

2026 Warble Court

Post Office Address

City

Thousand Oaks

State

CA

ZIP

91320

Country

U.S.A.

Burden Hour Statement: This form is estimated to take 0.4 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time you are required to complete this form should be sent to the Chief Information Officer, Patent and Trademark Office, Washington, DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Assistant Commissioner for Patents, Washington, DC 20231.

Please type a plus sign (+) inside this box → ☐

PTO/SB/02C (3-97)

Approved for use through 9/30/98. OMB 0651-0032

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number.

Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE

**DECLARATION****REGISTERED PRACTITIONER  
INFORMATION  
(Supplemental Sheet)**

Name	Registration Number	Name	Registration Number
Margaret M. Albanese	<u>45525</u>	Susan Lee	<u>30653</u>
Nancy V. Connelly	<u>46638</u>	Anita W. Magatti	<u>29825</u>
Eric S. Dicker	<u>31699</u>	Arthur Mann	<u>35598</u>
Robert A. Franks	<u>28605</u>	Christine F. Martin	<u>39762</u>
James M. Gould	<u>33702</u>	Edward H. Mazer	<u>27573</u>
Richard J. Grochala	<u>31518</u>	Jaye P. McLaughlin	<u>41211</u>
Henry S. Hadad	<u>35888</u>	Richard B. Murphy	<u>35296</u>
Thomas D. Hoffman	<u>28221</u>	James R. Nelson	<u>27929</u>
Henry C. Jeanette	<u>30856</u>	Pamela G. Salkeld	<u>38607</u>
Palaiyur S. Kalyanaraman	<u>34634</u>	David B. Schram	<u>43096</u>
Gerald P. Keleher	<u>43707</u>	Immac J. Thampoe	<u>36322</u>
Allan N. Kutzeno	<u>38945</u>	Paul A. Thompson	<u>35385</u>
		Donald W. Wyatt	<u>40879</u>
		Sandy Zaradic	<u>45997</u>

Burden Hour Statement: This form is estimated to take 0.4 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time you are required to complete this form should be sent to the Chief Information Officer, Patent and Trademark Office, Washington, DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Assistant Commissioner for Patents, Washington, DC 20231.